

QUANTUM HAMILTONIAN REDUCTION

YI SUN

1. THE QUANTIZATION FORMALISM

1.1. Preliminaries and notation. Throughout, we work over \mathbb{C} . Assume that \mathfrak{g} is the reductive Lie algebra associated to a reductive algebraic group G .

1.2. Three notions of quantization. Throughout this section, let A_0 be a commutative Poisson algebra. We describe the concepts of formal, graded, and filtered quantization of A_0 and how to translate between them under different conditions.

- (a) A *formal quantization* of A_0 is a $\mathbb{C}[[\hbar]]$ -algebra A , flat over $\mathbb{C}[[\hbar]]$ and complete and separated in the \hbar -adic topology, so that $A/\hbar A \simeq A_0$ and $\{a, b\} = \frac{1}{\hbar}[\bar{a}, \bar{b}] \pmod{\hbar}$. In this case, we also say that $(A, \{-, -\})$ is the *quasi-classical limit* of A .
- (b) Suppose that A_0 is $\mathbb{Z}_{\geq 0}$ -graded so that the Poisson bracket has degree -1 . A *graded quantization* (also called *algebraic quantization*) of A_0 over $\mathbb{C}[[\hbar]]$ is a $\mathbb{Z}_{\geq 0}$ -graded $\mathbb{C}[[\hbar]]$ -algebra A_{\hbar} so that $\deg(\hbar) = 1$ which is free as a $\mathbb{C}[[\hbar]]$ -module together with an isomorphism of graded algebras $A_{\hbar}/\hbar A_{\hbar} \rightarrow A_0$ such that $\{a, b\} = \frac{1}{\hbar}[\bar{a}, \bar{b}] \pmod{\hbar}$.
- (c) Suppose that A_0 is $\mathbb{Z}_{\geq 0}$ -graded so that the Poisson bracket has degree -1 . A *filtered quantization* of A_0 is a filtered algebra A where $[-, -]$ decreases the degree by 1 so that $\text{gr}A = A_0$ and $\text{gr}([\bar{a}, \bar{b}]) = \text{gr}(\{a, b\})$.

Suppose further that A_0 is $\mathbb{Z}_{\geq 0}$ -graded and that $\deg(\{-, -\}) = -1$. Given a formal quantization A_{\hbar} of A_0 equipped with a \mathbb{C}^* -action on A_{\hbar} inducing the $\mathbb{Z}_{\geq 0}$ -grading on A_0 , the subalgebra $A_{\hbar, \text{fin}} \subset A_{\hbar}$ of vectors with locally finite action of \mathbb{C}^* is a graded quantization of A_0 . Conversely, given a graded quantization $A_{\hbar, \text{fin}}$, we may form the corresponding formal quantization $A_{\hbar} := \widehat{A}_{\hbar, \text{fin}}$ as its \hbar -adic completion.

By Propositions 1.1 and 1.2, we may convert between filtered and graded quantizations. For a filtered quantization A_f of A_0 , the construction proceeds via the Rees algebra

$$\text{Rees}(A_f) = \bigoplus_{n \geq 0} A_f^n \cdot \hbar^n \subset A_f[[\hbar]].$$

Proposition 1.1. If A_f is a filtered quantization of A_0 , $A_{\hbar} := \text{Rees}(A_f)$ viewed as a graded $\mathbb{C}[[\hbar]]$ -subalgebra of $A_f[[\hbar]]$, is a graded quantization of A_0 .

Proof. By construction, we see that $A_{\hbar}/\hbar A_{\hbar} \simeq \text{gr}A_f = A_0$. Further, for elements $a^i \in A_f^i$ and $a^j \in A_f^j$, because A_f is a filtered quantization of A_0 , we have $[a^i, a^j] \in A_f^{i+j-1}$ and $\text{gr}([a^i, a^j]) = \text{gr}(\{a^i, a^j\})$. Therefore, in A_{\hbar} , we have $[a^i, a^j] \in \hbar A_{\hbar}^{i+j-1}$, so that $\frac{1}{\hbar}[a^i, a^j] = \{a^i, a^j\} \pmod{\hbar}$. \square

Proposition 1.2. Suppose that A_{\hbar} is a graded quantization of A_0 which is $\mathbb{Z}_{\geq 0}$ -graded so that $\deg(\hbar) = 1$. Then $A_f := A_{\hbar}/(\hbar - 1)A_{\hbar}$ with filtration induced from the grading of A_{\hbar} is a filtered quantization of A_0 .

Proof. Denote the graded parts of A_{\hbar} by A_{\hbar}^n and the corresponding filtration by $A_{\hbar}^{\leq n}$. Because the element \hbar has degree 1, we find that

$$\text{gr}A_{\hbar} = \bigoplus_{n \geq 0} (A_{\hbar}^{\leq n} / A_{\hbar}^{\leq n-1}) / (\hbar - 1)(A_{\hbar}^{\leq n-1} / A_{\hbar}^{\leq n-2}) = \bigoplus_{n \geq 0} A_{\hbar}^n / \hbar A_{\hbar}^{n-1} = A_{\hbar} / \hbar A_{\hbar}.$$

Further, for $a^i \in A_{\hbar}^i$ and $a^j \in A_{\hbar}^j$, we see that

$$[a^i, a^j] = \{a^i, a^j\} \pmod{\hbar} \text{ in } A_{\hbar}^{i+j} / (\hbar - 1)A_{\hbar}^{i+j-1},$$

which shows exactly that $\text{gr}([a^i, a^j]) = \text{gr}(\{a^i, a^j\})$. \square

1.3. Two homogenized algebras. We now give two examples of graded quantizations which will be important in the next section. Define the *homogenized universal enveloping algebra* of a Lie algebra \mathfrak{g} to be the $\mathbb{C}[\hbar]$ -algebra

$$U_{\hbar}(\mathfrak{g}) := \left\langle \hbar, x \in \mathfrak{g} \mid x \star y - y \star x = \hbar[x, y] \right\rangle,$$

where \star denotes the product in $U_{\hbar}(\mathfrak{g})$. Notice that $U_{\hbar}(\mathfrak{g})$ is a graded quantization of $S(\mathfrak{g})$ and that the corresponding filtered quantization $U_{\hbar}(\mathfrak{g})/(\hbar-1)U_{\hbar}(\mathfrak{g}) \simeq U(\mathfrak{g})$ is the standard universal enveloping algebra. The corresponding Poisson algebra structure on $S(\mathfrak{g})$ is the one induced by $\{x, y\} = [x, y]$.

Let X be a smooth affine algebraic variety. Define the ring of *homogenized differential operators on X* to be the $\mathbb{C}[\hbar]$ -algebra generated by \mathcal{O}_X in grade 0, $\text{Vect}(X)$ in grade 1 and subject to the relations

$$D_{\hbar}(X) := \langle f, v \mid f \star g = f \cdot g, f \star v = f \cdot v, v \star f = f \cdot v + \hbar v(f), u \star v - v \star u = \hbar[u, v] \rangle,$$

where we use \star to denote the product in $D_{\hbar}(X)$. Observe that $D_{\hbar}(X)$ is a graded quantization of $p_*\mathcal{O}_{T^*X}$ (where $p : T^*X \rightarrow X$ is the natural projection) and that $D_{\hbar}(X)/(\hbar-1)D_{\hbar}(X) = D(X)$, the ordinary ring of differential operators on X .

2. QUANTUM REDUCTION FORMALISM

We now define quantum reduction, first in the general case and then in the case of a graded quantization.

2.1. Quantum moment map. Let A be an associative algebra equipped with a locally-finite completely reducible \mathfrak{g} -action given by a map of Lie algebras $\phi : \mathfrak{g} \rightarrow \text{Der}(A)$. We say that $\mu : U(\mathfrak{g}) \rightarrow A$ is a *quantum moment map* for ϕ if $\phi(x)(-) = [\mu(x), -]$ so that the action of \mathfrak{g} factors through μ .

Example. Let X be an affine algebraic variety with an action of G . Differentiation of the group action gives rise to a Lie algebra map $\mu : \mathfrak{g} \rightarrow D(X)$, where $D(X)$ is the algebra of differential operators on X .

In the homogenized context, if A_{\hbar} is a $\mathbb{Z}_{\geq 0}$ -graded $\mathbb{C}[\hbar]$ -algebra so that $\deg(\hbar) = 1$ with grading-preserving \mathfrak{g} -action $\phi : U_{\hbar}(\mathfrak{g}) \rightarrow \text{Der}(A_{\hbar})$, we say that $\mu_{\hbar} : U_{\hbar}(\mathfrak{g}) \rightarrow A_{\hbar}$ is a *homogenized quantum moment map* if the action is given by $\phi(x) = \frac{1}{\hbar}[\mu_{\hbar}(x), -]$ and $\mu_{\hbar}(\mathfrak{g}) \subset A_{\hbar}^1$. In the previous example, we may obtain a *homogenized quantum moment map* $\mu_{\hbar} : U_{\hbar}(\mathfrak{g}) \rightarrow D_{\hbar}(X)$ in the same way. Observe that $\mu = \mu_{\hbar} \pmod{\hbar-1}$.

2.2. Definition of quantum reduction. Suppose now that we have either a quantum moment map μ or homogenized quantum moment map μ_{\hbar} . For a character $\lambda : \mathfrak{g} \rightarrow \mathbb{C}$, let $J_{\mu, \lambda} := A \cdot \text{span}\{\mu(x) - \langle \lambda, x \rangle \mid x \in \mathfrak{g}\}$ (in the graded context $J_{\mu, \lambda, \hbar} := A_{\hbar} \cdot \text{span}\{\mu_{\hbar}(x) - \hbar \langle \lambda, x \rangle \mid x \in \mathfrak{g}\}$, a graded ideal). Observe that the \mathfrak{g} -action fixes $J_{\mu, \lambda}$ and $J_{\mu, \lambda, \hbar}$. We define the *quantum Hamiltonian reduction* of A with respect to $\lambda : \mathfrak{g} \rightarrow \mathbb{C}$ and μ to be the algebra

$$R(\mathfrak{g}, A, \lambda) := \text{End}_{\mathfrak{g}}(A/J_{\mu, \lambda}) \simeq (A/J_{\mu, \lambda})^{\mathfrak{g}} \simeq A^{\mathfrak{g}}/J_{\mu, \lambda}^{\mathfrak{g}},$$

where the algebra structure is by composition in the first expression and induced by multiplication in A and $A^{\mathfrak{g}}$ in the last two expressions. In particular, we see that $(a_1 + J_{\mu, \lambda}) \cdot (a_2 + J_{\mu, \lambda}) = a_1 a_2 + J_{\mu, \lambda}$.

Remark. We make several comments on this definition.

- The first two expressions for $R(\mathfrak{g}, A, \lambda)$ may be defined for any Lie algebra \mathfrak{g} and are isomorphic without any assumption on \mathfrak{g} and μ .
- The second isomorphism sends $\phi \in \text{End}_{\mathfrak{g}}(A/J_{\mu, \lambda})$ to $\phi(1) \in (A/J_{\mu, \lambda})^{\mathfrak{g}}$, where we note that $\phi(1)$ is \mathfrak{g} -invariant because 1 is.
- In general, $J_{\mu, \lambda}$ is not a two-sided ideal in A , so $A/J_{\mu, \lambda}$ does not carry an algebra structure.
- The final isomorphism relies on the fact that \mathfrak{g} is a reductive Lie algebra acting locally finitely and completely reducibly on A . In this case, Lemma 2.1 shows that $J_{\mu, \lambda}$ becomes a two-sided ideal after taking \mathfrak{g} -invariants, so we may consider the final expression as a quotient algebra of $A^{\mathfrak{g}}$.

Lemma 2.1. The invariant ideal $J_{\mu, \lambda}^{\mathfrak{g}}$ is a two-sided ideal in $A^{\mathfrak{g}}$.

Proof. For any $b \in A^{\mathfrak{g}}$, we have $[\mu(x) - \langle \lambda, x \rangle, b] = 0$ by definition, hence for any element $\xi = a(\mu(x) - \langle \lambda, x \rangle) \in J_{\mu, \lambda}^{\mathfrak{g}}$, we see that $\xi b = ab(\mu(x) - \langle \lambda, x \rangle) \in J_{\mu, \lambda}^{\mathfrak{g}}$. \square

If A_{\hbar} is a $\mathbb{Z}_{\geq 0}$ -graded $\mathbb{C}[\hbar]$ -algebra, we may define the homogenized reduction $R_{\hbar}(\mathfrak{g}, A_{\hbar}, \lambda)$ in the same way by replacing $J_{\mu, \lambda}$ by $J_{\mu, \lambda, \hbar}$ in the discussion above. Note that $R_{\hbar}(\mathfrak{g}, A_{\hbar}, \lambda)$ is $\mathbb{Z}_{\geq 0}$ -graded because the \mathfrak{g} -action preserves grading and $J_{\mu_{\hbar}, \lambda, \hbar}^{\mathfrak{g}}$ is a graded ideal.

2.3. Universal reduction. For the rest of the talk, it will be useful to consider the *universal reduction* of A with respect to μ , which is the algebra defined by

$$R(\mathfrak{g}, A) := A^{\mathfrak{g}} / (A\mu([\mathfrak{g}, \mathfrak{g}]))^{\mathfrak{g}}.$$

Multiplying on the left via $\mu : U(\mathfrak{g}) \rightarrow A$ and projecting to \mathfrak{g} -invariants gives a $U(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])$ -action on $R(\mathfrak{g}, A)$, where we note that $U(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]) \simeq S(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])$. Further, we see that

$$R(\mathfrak{g}, A, \lambda) = \mathbb{C}_\lambda \otimes_{S(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])} R(\mathfrak{g}, A),$$

where \mathbb{C}_λ is the $S(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])$ -module associated with the character λ . In a similar way, we may define $R_{\hbar}(\mathfrak{g}, A_{\hbar})$, which is a $S_{\hbar}(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])$ -algebra so that

$$R_{\hbar}(\mathfrak{g}, A_{\hbar}, \lambda) = \mathbb{C}_\lambda[\hbar] \otimes_{S_{\hbar}(\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}])} R_{\hbar}(\mathfrak{g}, A_{\hbar}).$$

3. QUANTIZATION COMMUTES WITH REDUCTION

3.1. Classical reduction. We now wish to view the quantum reduction as a quantization of the classical Hamiltonian reduction. Let (M, ω) be a symplectic algebraic variety with G -action which induces a map of Lie algebras $\phi_0 : \mathfrak{g} \rightarrow \text{Der}(\mathbb{C}[M])$. A *classical co-moment map* μ_0 may be viewed as a map of Poisson algebras

$$\mu_0 : S(\mathfrak{g}) \rightarrow \mathbb{C}[M]$$

so that $\{\mu_0(x), f\} = \phi_0(x) \cdot f$. Define the left Poisson ideal

$$I_{\mu_0} := \mathbb{C}[M] \cdot \mu_0(\mathfrak{g})$$

Recall that the *classical Hamiltonian reduction* of M with respect to λ is

$$R(G, M, 0) := (\mathbb{C}[M]/I_{\mu_0})^G \simeq \mathbb{C}[M]^G / I_{\mu_0}^G.$$

Geometrically, it corresponds to the space $(\mu_0^*)^{-1}(0)//G = \text{Spec}(R(G, M, 0))$.

3.2. Quantum reduction as a quantization. We now relate the quantum and classical reductions. We will consider mainly the graded context, but will provide the corresponding statements in the filtered context. Let A_{\hbar} be a graded quantization of $A_0 := \mathbb{C}[M]$. If $\mu_{\hbar} : U_{\hbar}(\mathfrak{g}) \rightarrow A_{\hbar}$ is a homogenized quantum moment map, then $\mu_0 := \mu \pmod{\hbar}$ is a classical co-moment map for the G -action on M . Under certain conditions, Proposition 3.1 allows us to relate the two types of reduction.

Proposition 3.1. Let $I_0 \subset A_0$ be the ideal generated by $\mu_0(\mathfrak{g})$. Let x_1, \dots, x_k be a basis for \mathfrak{g} . If $\mu_0(x_1), \dots, \mu_0(x_k)$ form a regular sequence in A_0 , then for any λ , the quantum reduction $R_{\hbar}(\mathfrak{g}, A_{\hbar}, \lambda)$ is a graded quantization of the classical reduction $R(G, M, 0)$.

Proof. Write $\mathfrak{z} = \mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ and consider the decomposition $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}] \oplus \mathfrak{z}$. Note that $\mu_0(x_1), \dots, \mu_0(x_k)$ is a regular sequence if and only if $\text{Spec}(A_0/I_0)$ has codimension k in $\text{Spec}(A_0)$, so we may assume the basis $\{x_i\}$ is chosen so that x_1, \dots, x_m form a basis of $[\mathfrak{g}, \mathfrak{g}]$ and x_{m+1}, \dots, x_k form a basis of \mathfrak{z} .

We first claim that $I = A_{\hbar}\mu([\mathfrak{g}, \mathfrak{g}])$ is saturated, meaning that $I \cap \hbar A_{\hbar} = \hbar I$. Suppose that $\hbar a \in I$ for some $a \in A_{\hbar}$; then we can write $\hbar a = \sum_i a_i \mu(x_i)$ for some $a_i \in A_{\hbar}$. Passing to the quotient, we see that $\sum_i a_i^0 \mu_0(x_i) = 0$, where a_i^0 denotes the image of a_i in A_0 . Because $\mu_0(x_i)$ form a regular sequence, the first homology of the associated Koszul complex

$$\bigoplus_{i < j} A_0 \xrightarrow{\sum_{i \neq j} a_{ij}^0 \mu_0(x_j)} \bigoplus_i A_0 \xrightarrow{\sum_i a_i^0 \mu_0(x_i)} A_0$$

is trivial, where we adopt the convention that $a_{ji}^0 = -a_{ij}^0$ for $i < j$. Therefore, we may write

$$a_i^0 = \sum_{i \neq j} a_{ij}^0 \mu_0(x_j)$$

for some a_{ij}^0 . We conclude that $a_i = \sum_{i \neq j} a_{ij} \mu(x_j) + \hbar b_i$ for lifts a_{ij} of a_{ij}^0 and $b_i \in A_{\hbar}$, hence

$$a = \sum_i \sum_{j \neq i} a_{ij} \mu(x_j) \mu(x_i) + \hbar \sum_i b_i \mu(x_i) = \sum_{i < j} a_{ij} [\mu(x_j), \mu(x_i)] + \hbar \sum_i b_i \mu(x_i) \in \hbar I,$$

since $[\mu(x_i), \mu(x_j)] = \hbar \mu([x_i, x_j])$. We obtain the desired $I \cap \hbar A_{\hbar} = \hbar I$.

We claim now that A_{\hbar}/I is graded free over $S_{\hbar}(\mathfrak{z})$. Saturation and the given regularity condition mean exactly that $\hbar, \mu_0(x_{m+1}), \dots, \mu_0(x_k)$ form a A_{\hbar}/I -regular sequence, so this follows from Lemma 3.2. We conclude further that $R_{\hbar}(\mathfrak{g}, A) = (A_{\hbar}/I)^{\mathfrak{g}}$ is graded free over $S_{\hbar}(\mathfrak{z})$ as well because the \mathfrak{g} -action commutes with the action of $S_{\hbar}(\mathfrak{z})$.

Lemma 3.2. Let M be a graded $k[x_1, \dots, x_n]$ -module so that x_1, \dots, x_n are a M -regular sequence and $\deg(x_i) = 1$. Then M is a (graded) free module over $k[x_1, \dots, x_n]$.

Proof. By induction on n , we may reduce to the case where $n = 1$, where the statement is obvious. \square

Observe now that $R_{\hbar}(\mathfrak{g}, A, \lambda) = \mathbb{C}_{\lambda}[\hbar] \otimes_{S_{\hbar}(\mathfrak{z})} R_{\hbar}(\mathfrak{g}, A)$ is evidently flat over $\mathbb{C}[\hbar]$ and that

$$\begin{aligned} R_{\hbar}(\mathfrak{g}, A, \lambda)/\hbar R_{\hbar}(\mathfrak{g}, A, \lambda) &= \mathbb{C} \otimes_{\mathbb{C}[\hbar]} \mathbb{C}_{\lambda}[\hbar] \otimes_{S_{\hbar}(\mathfrak{z})} R_{\hbar}(\mathfrak{g}, A) \\ &= \mathbb{C} \otimes_{S(\mathfrak{z})} S(\mathfrak{z}) \otimes_{S_{\hbar}(\mathfrak{z})} R_{\hbar}(\mathfrak{g}, A) \\ &= \mathbb{C} \otimes_R (\mathfrak{g}, A_0) \\ &= R(G, M, 0). \end{aligned}$$

We conclude that $R_{\hbar}(\mathfrak{g}, A, \lambda)$ is a graded quantization of $R(G, M, 0)$ for all λ . \square

Consider now the corresponding filtered setting. Let A be a filtered quantization of A_0 and $\mu : U(\mathfrak{g}) \rightarrow A$ a quantum moment map so that $\mu_0 := \text{gr}(\mu)$ is a classical co-moment map. Translating Proposition 3.1 into this filtered setting, we obtain Corollary 3.3.

Corollary 3.3. Let $I_0 \subset A_0$ be the ideal generated by $\mu_0(\mathfrak{g})$. Let x_1, \dots, x_k be a basis for \mathfrak{g} . If $\mu_0(x_1), \dots, \mu_0(x_k)$ form a regular sequence in A_0 , then for any λ , the quantum reduction $R(\mathfrak{g}, A, \lambda)$ is a filtered quantization of the classical reduction $R(G, M, 0)$.

Proof. Each of the constructions in the filtered case is obtained by reducing the corresponding construction in the graded case modulo the ideal generated by $(\hbar - 1)$, so this follows from Proposition 3.1. \square

REFERENCES

- [1] P. Etingof, Lectures on Calogero-Moser systems (2005). Available at: <http://arxiv.org/pdf/math/0606233v4.pdf>
- [2] P. Etingof and V. Ginzburg, Symplectic reflection algebras, Calogero-Moser space, and deformed Harish-Chandra homomorphism, *Inventiones mathematicae* **147** (2002), pp. 243-348. Available at: <http://arxiv.org/abs/math/0011114>.
- [3] W. L. Gan and V. Ginzburg, Almost-commuting variety, D -modules, and Cherednik algebras, *International Mathematics Research Papers* **2006** (2005). Article ID 26439, pp 154. Available at: <http://arxiv.org/abs/math/0409262>.
- [4] I. Losev, Isomorphisms of quantizations via quantization of resolutions, *Advances in Mathematics* **231** (2012), pp. 1216-1270. Available at: <http://arxiv.org/abs/1010.3182>.
- [5] I. Losev, Lectures on symplectic reflection algebras (2012). Available at: http://www.northeastern.edu/iloseu/SRA/SRA_lec14.pdf.

E-mail address: yisun@math.mit.edu