

# 54th International Mathematical Olympiad

JOHN BERMAN  
University of Virginia  
Charlottesville, VA 22903  
jdb8pc@virginia.edu

ZUMING FENG  
Phillips Exeter Academy  
Exeter, NH 03833  
zfeng@exeter.edu

YI SUN  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
yisun@mit.edu

## Problems (Day 1)

1. Prove that for any pair of positive integers  $k$  and  $n$ , there exist  $k$  positive integers  $m_1, m_2, \dots, m_k$  (not necessarily different) such that

$$1 + \frac{2^k - 1}{n} = \left(1 + \frac{1}{m_1}\right) \left(1 + \frac{1}{m_2}\right) \cdots \left(1 + \frac{1}{m_k}\right).$$

2. A configuration of 4027 points in the plane is called *Colombian* if it consists of 2013 red points and 2014 blue points, and no three of the points of the configuration are collinear. By drawing some lines, the plane is divided into several regions. An arrangement of lines is *good* for a Colombian configuration if the following two conditions are satisfied:
- no line passes through any point of the configuration;
  - no region contains points of both colours.

Find the least value of  $k$  such that for any Colombian configuration of 4027 points, there is a good arrangement of  $k$  lines.

3. Let the excircle of the triangle  $ABC$  opposite the vertex  $A$  be tangent to side  $BC$  at  $A_1$ . Define the points  $B_1$  on  $CA$  and  $C_1$  on  $AB$  analogously, using the excircles opposite  $B$  and  $C$ , respectively. Suppose that the circumcentre of triangle  $A_1B_1C_1$  lies on the circumcircle of triangle  $ABC$ . Prove that triangle  $ABC$  is right-angled.

*The excircle of triangle  $ABC$  opposite the vertex  $A$  is the circle that is tangent to the line segment  $BC$ , to the ray  $AB$  beyond  $B$ , and to the ray  $AC$  beyond  $C$ . The excircles opposite  $B$  and  $C$  are similarly defined.*

## Problems (Day 2)

4. Let  $ABC$  be an acute-angled triangle with orthocenter  $H$ , and let  $W$  be a point on the side  $BC$ , lying strictly between  $B$  and  $C$ . The points  $M$  and  $N$  are the feet of the altitudes from  $B$  and  $C$ , respectively. Denote by  $\omega_1$  the circumcircle of  $BWN$ , and let  $X$  be the point on  $\omega_1$  such that  $WX$  is a diameter of  $\omega_1$ . Analogously, denote by  $\omega_2$  the circumcircle of triangle  $CWM$ , and let  $Y$  be the point on  $\omega_2$  such that  $WY$  is a diameter of  $\omega_2$ . Prove that  $X$ ,  $Y$ , and  $H$  are collinear.

5. Let  $\mathbb{Q}_{>0}$  be the set of positive rational numbers. Let  $f : \mathbb{Q}_{>0} \rightarrow \mathbb{R}$  be a function satisfying the following three conditions:

- (i) for all  $x, y \in \mathbb{Q}_{>0}$ , we have  $f(x)f(y) \geq f(xy)$ ;
- (ii) for all  $x, y \in \mathbb{Q}_{>0}$ , we have  $f(x + y) \geq f(x) + f(y)$ ;
- (iii) there exists a rational number  $a > 1$  such that  $f(a) = a$ .

Prove that  $f(x) = x$  for all  $x \in \mathbb{Q}_{>0}$ .

6. Let  $n \geq 3$  be an integer, and consider a circle with  $n + 1$  equally spaced points marked on it. Consider all labellings of these points with the numbers  $0, 1, \dots, n$  such that each label is used exactly once; two such labellings are considered to be the same if one can be obtained from the other by a rotation of the circle. A labelling is called *beautiful* if, for any four labels  $a < b < c < d$  with  $a + d = b + c$ , the chord joining the points labeled  $a$  and  $d$  does not intersect the chord joining the points labeled  $b$  and  $c$ .

Let  $M$  be the number of beautiful labellings, and let  $N$  be the number of ordered pairs  $(x, y)$  of positive integers such that  $x + y \leq n$  and  $\gcd(x, y) = 1$ . Prove that

$$M = N + 1.$$

### Solutions

1. We induct on  $k$ . The base case  $k = 1$  holds with  $m_1 = n$ . If the statement holds for some  $k$ , we consider two cases. If  $n = 2m - 1$  is odd, we have

$$1 + \frac{2^{k+1} - 1}{n} = \frac{2m}{2m - 1} \frac{2^{k+1} + 2m - 2}{2m} = \left(1 + \frac{1}{2m - 1}\right) \left(1 + \frac{2^k - 1}{m}\right),$$

where  $1 + \frac{2^k - 1}{m}$  is the product of  $k$  terms of the desired form by the induction hypothesis, yielding the desired decomposition. If  $n = 2m$  is even, we have

$$\begin{aligned} 1 + \frac{2^{k+1} - 1}{n} &= \frac{2^{k+1} + 2m - 1}{2^{k+1} + 2m - 2} \frac{2^{k+1} + 2m - 2}{2m} \\ &= \left(1 + \frac{1}{2^{k+1} + 2m - 2}\right) \left(1 + \frac{2^k - 1}{m}\right), \end{aligned}$$

where  $1 + \frac{2^k - 1}{m}$  is the product of  $k$  terms of the desired form by the induction hypothesis, yielding the desired decomposition and completing the induction.

This problem was proposed by the Olympiad problem committee from Japan.

2. The answer is 2013. We first show that a good arrangement with at most 2013 lines always exists. We begin with the following key lemma.

**LEMMA 1.** *Any pair of points  $P$  and  $Q$  in a Colombian configuration  $C$  can be separated from the other points by two lines.*

*Proof.* No three points in  $C$  are collinear, so each other point has one of finitely many positive distances to  $PQ$ . Choose  $r > 0$  less than all such distances; the two lines parallel to and at a distance  $r$  from  $PQ$  have the desired property. ■

Let  $\mathcal{C}$  be the convex hull of our Colombian configuration  $C$ . If a red point  $R$  is a vertex of  $\mathcal{C}$ , draw a line  $\ell_1$  separating  $R$  from all other points. Next, place the other 2012 red points into 1006 pairs and apply Lemma 1 to draw 2012 lines separating them from the rest of  $C$ . Together with  $\ell_1$ , these 2012 lines form a good arrangement. Otherwise,  $\mathcal{C}$  has a side  $B_1B_2$  consisting of blue points. Draw a line  $\ell_1$  separating  $B_1$  and  $B_2$  from the rest of  $C$ , place the 2012 remaining blue points into

1006 pairs, and apply Lemma 1. The resulting 2012 lines together with  $\ell_1$  form the desired good arrangement.

For the other direction, let  $\mathcal{P} = A_1 \cdots A_{4026}$  be a regular 4026-gon with  $A_i$  red for  $i$  odd and blue for  $i$  even. Form a Colombian configuration  $C$  with the vertices of  $\mathcal{P}$  and another blue point  $B$ . We claim that any good arrangement for  $C$  has at least 2013 lines. Note that each of the 4026 pairs  $(A_i, A_{i+1})$  of neighboring points contains points in different regions, so  $A_i A_{i+1}$  intersects at least one of the lines. Each line intersects  $\mathcal{P}$  at most twice, so an arrangement with  $k$  lines can produce at most  $2k$  intersections, implying that  $2k \geq 4026$  and hence  $k \geq 2013$ , as desired.

This problem was proposed by Ivan Guo from Australia. The current formulation of this problem was suggested during IMO jury meetings by Leonardo I. M. Sandoval from Mexico.

3. Let  $\omega$  be the circumcircle of  $ABC$ , and let  $O_1$  be the circumcenter of  $A_1 B_1 C_1$ . Because  $A_1, B_1,$  and  $C_1$  are on the boundary of  $ABC$  and  $O_1$  is outside of  $ABC$ ,  $A_1 B_1 C_1$  is obtuse. Without loss of generality, assume that  $\angle B_1 A_1 C_1$  is obtuse so that  $O_1$  and  $A$  lie on the same side of line  $B_1 C_1$ .

LEMMA 2. *The second intersection  $A_0$  of  $\omega$  and the circumcircle of triangle  $AB_1 C_1$  is the midpoint of arc  $\widehat{BAC}$ .*

*Proof.* By the definition of  $A_0$ , we have  $\angle A_0 B C_1 = \angle A_0 B A = \angle A_0 C A = \angle A_0 C B_1$  and  $\angle A_0 C_1 A = \angle A_0 B_1 A$ ; hence,  $AC_1 B$  and  $AB_1 C$  are similar. But  $BC_1 = CB_1$ , so these two triangles are congruent; hence,  $A_0 B = A_0 C$ . Because  $AA_0 B_1 C_1$  is cyclic, we have  $\angle C_1 A_0 B_1 = \angle C_1 A B_1 = \angle BAC$ , so  $A_0$  lies on  $\widehat{BAC}$  with  $BA_0 = CA_0$ , implying that  $A_0$  is the midpoint of  $\widehat{BAC}$ . ■

By Lemma 2, a spiral similarity centered at  $A_0$  sends  $B_1 C_1$  to  $CB$ , so  $A_0$  is the intersection of  $\omega$  and the perpendicular bisector of  $B_1 C_1$ , which is on the same side of  $BC$  as  $A$ . Recalling that  $A_0$  is the circumcenter of  $A_1 B_1 C_1$ , and using this result for the analogous points  $B_0$  and  $C_0$ , we obtain that  $A_0 C_1 B_0 A_1$  and  $A_0 A_1 C_0 B_1$  are kites with symmetry axes  $A_0 B_0$  and  $A_0 C_0$ . Recalling that  $C_1 B_1 A A_0$  is cyclic, we have  $\angle CAB = \angle C_1 A_0 B_1 = 2\angle B_0 A_0 C_0 = \widehat{B_0 C_0}$ . By Lemma 2,  $B_0$  and  $C_0$  are the midpoints of  $\widehat{ABC}$  and  $\widehat{BCA}$ , hence

$$\begin{aligned} \angle CAB &= \widehat{B_0 C_0} = 360^\circ - \widehat{ACC_0} - \widehat{B_0 A} = 360^\circ - \frac{\widehat{BCA} + \widehat{ABC}}{2} \\ &= 360^\circ - \frac{360^\circ - 2\angle BCA + 360^\circ - 2\angle ABC}{2} = \angle BCA + \angle ABC, \end{aligned}$$

implying that  $\angle CAB = 90^\circ$ , so  $ABC$  has right angle at vertex  $A$ .

This problem was proposed by Alexander Polyanskiy from Russia.

4. Let  $L$  be the foot of the altitude of  $ABC$  from  $A$ , and let  $O_1$  and  $O_2$  be the centers of  $\omega_1$  and  $\omega_2$ , respectively. Because  $\angle WNB < \angle CNB = 90^\circ$ ,  $O_1$  and  $N$  lie on the same side of  $BC$ . Likewise,  $O_2$  and  $M$  lie on the same side of  $BC$ . Hence, segment  $O_1 O_2$  does not intersect line  $BC$ . In particular,  $W$  does not lie on line  $O_1 O_2$ , and  $\omega_1$  and  $\omega_2$  intersect again at a point  $Z$  other than  $W$ .

Because  $XW$  is a diameter of  $\omega_1$ , we have  $XZ \perp WZ$ . Likewise, we have  $YZ \perp WZ$ , so  $X, Y,$  and  $Z$  lie on a line perpendicular to line  $ZW$ . It suffices to show that  $HZ \perp ZW$ . First, quadrilaterals  $BNHL$  and  $CMHL$  are cyclic, so by power of a point,  $AM \cdot AC = AH \cdot AL = AN \cdot AB$ ; hence,  $A$  lies on the radical axis  $ZW$  of  $\omega_1$  and  $\omega_2$ . Second, by our previous argument,  $A$  is the radical center of circles  $\omega_1, \omega_2$ , and the circumcircles of quadrilaterals  $BNHL$  and  $CMHL$ . In particular, we have

$AH \cdot AL = AZ \cdot AW$ , so  $ZHLW$  is cyclic. This shows that  $\angle AZH = \angle ALW = 90^\circ$ , hence  $HZ \perp AW$ , completing the proof.

This problem was proposed by Warut Suksompong and Potcharapol Suteparuk from Thailand.

5. We claim that the only solution is  $f(x) = x$  for all  $x \in \mathbb{Q}_{>0}$ .

By applying (i) and iterating (ii), we see that  $f(n)f(x) \geq f(nx) \geq nf(x)$  for positive integer  $n$ . Setting  $x = a$ , we see that  $f(n) \geq n$ , because  $f(a) = a > 0$ .

We claim that  $f$  is non-negative and non-decreasing. Indeed, if  $f(y) < 0$ , setting  $x = y$  and dividing by  $f(y)$  in our first inequality shows that  $f(n) \leq n$ , hence  $f(n) = n$ . This chain of inequalities is thus an equality, so  $f(n)f(x) = f(nx)$  for all  $x$ . Writing  $y = \frac{p}{q} \in \mathbb{Q}_{>0}$ , we have  $f(q)f(\frac{p}{q}) = f(p)$ , so  $f(y) = y$ , a contradiction. Hence,  $f$  is non-negative, thus also non-decreasing by (ii).

We claim now that  $f(x) \geq x$  for all  $x \geq 1$ . First, note that  $f(x) \geq f(\lfloor x \rfloor) \geq \lfloor x \rfloor > x - 1$ . From (i) we know that  $f(x)^n \geq f(x^n)$ , so  $f(x)^n \geq f(x^n) > x^n - 1$ . But if  $f(x) = x - \epsilon$  for some  $\epsilon > 0$  and  $x > 1$ , then for all  $n$  we have  $1 > x^n - f(x)^n \geq (x - f(x))(x^{n-1}) = \epsilon x^{n-1}$ . Since  $x > 1$ , we can choose  $n$  such that  $x^{n-1} > \frac{1}{\epsilon}$ , a contradiction. Therefore,  $f(x) \geq x$  for all  $x > 1$ , and we already know that  $f(1) \geq 1$ , yielding the claim.

We now show that  $f(x) = x$  for  $x \geq 1$ . Note that  $a^k = f(a)^k \geq f(a^k)$  for positive integers  $k$  by (i). We also have  $f(a^k) \geq a^k$ , so  $f(a^k) = a^k$  for positive integers  $k$ . For  $x \geq 1$  and  $k$  with  $a^k > 2x$ , we have  $a^k = f(a^k) \geq f(x) + f(a^k - x) \geq x + (a^k - x) = a^k$ . Equality thus holds, so  $f(x) = x$  for  $x \geq 1$ .

Finally, for any integer  $n$ , we have  $f(n) = n$  and  $f(n)f(x) \geq f(nx) \geq nf(x)$ , so equality holds, implying that  $f(nx) = nf(x)$ . In particular, for any  $x = \frac{p}{q}$  in  $\mathbb{Q}_{>0}$ , we conclude that  $qf(x) = f(p) = p$ , hence  $f(x) = x$ , as desired.

This problem was proposed by Nikolai Nikolov from Bulgaria.

6. We will prove that there are  $N + 1$  beautiful labelings for all  $n \geq 2$ .

Let  $0 < x < 1$  be a real number. Define the beautiful labeling  $C_n(x)$  as follows. For  $0 \leq k \leq n$ , let  $W_k = e^{\frac{2\pi i k}{n}} x$ , and let  $Z_k$  be the point that results from rearranging the  $W_k$  evenly in the same relative position; label  $Z_k$  by  $k$ . Note that  $W_a W_b$  and  $W_c W_d$  intersect iff  $Z_a Z_b$  and  $Z_c Z_d$  intersect.

Call such a labeling *cyclic*, and call it *degenerate* if two of the  $W_k$  coincide. Note that  $C_n(x)$  is degenerate iff  $x$  is a reduced fraction with denominator at most  $n$ . Call such numbers *good*; there are  $N$  good fractions in  $(0, 1)$ .

LEMMA 3. *A labeling is beautiful if and only if it is non-degenerate cyclic.*

*Proof.* Let  $C_n(x)$  be a non-degenerate cyclic labeling. For any  $0 \leq a < b < c < d \leq n$  with  $a + d = b + c$ , arcs  $\widehat{W_a W_b}$  and  $\widehat{W_c W_d}$  have the same measure, so  $W_a W_d \parallel W_b W_c$ , implying that  $C_n(x)$  is beautiful.

For the converse, induct on  $n$  with trivial base case  $n = 2$ . If all beautiful arrangements of  $[0, n - 1]$  are cyclic, for a beautiful arrangement  $A$  of  $[0, n]$ , form  $A' = C_{n-1}(x)$  by removing  $n$ . Let  $x$  lie between consecutive good fractions  $p_1/q_1$  and  $p_2/q_2$  with  $q_1, q_2 \leq n - 1$ , and consider two cases.

*Case 1:* There is no fraction with denominator  $n$  between  $\frac{p_1}{q_1}$  and  $\frac{p_2}{q_2}$ . If  $A \neq C_n(x)$ , they can differ only in the location of  $n$ . Suppose  $W_n$  occurs directly between  $W_i$  and  $W_j$  in  $C_n(x)$  in clockwise order. Note that  $i + (n - 1) = (i - 1) + n$  and  $j + (n - 1) = (j - 1) + n$ , so the two corresponding pairs of chords do not intersect and  $W_i, W_n, W_j, W_{i-1}, W_{n-1}, W_{j-1}$  occur in that order. In  $A$ , since  $W_i W_{n-1}$  does not intersect  $W_n W_{i-1}$ ,  $W_n$  lies on arc  $\widehat{W_i W_{n-1}}$ . Likewise,  $W_n$  must lie on arc  $\widehat{W_{n-1} W_j}$ . Thus  $W_n$  lies between  $W_i$  and  $W_j$ , so  $A = C_n(x)$ .

*Case 2:* There is a fraction  $\frac{a}{n}$  with denominator  $n$  between  $\frac{p_1}{q_1}$  and  $\frac{p_2}{q_2}$ . Since  $\frac{p_2}{q_2} -$

$\frac{p_1}{q_1} \leq \frac{1}{n-1}$ , there is a unique such fraction. Choose  $x_1 \in (\frac{p_1}{q_1}, \frac{a}{n})$  and  $x_2 \in (\frac{a}{n}, \frac{p_2}{q_2})$ . We wish to show that either  $A = C_n(x_1)$  or  $A = C_n(x_2)$ . In  $A'$ ,  $W_{q_1}, W_0, W_{q_2}$  occur in that clockwise order. It suffices to show that  $W_n$  lies on arc  $\widehat{W_{q_1} W_{q_2}}$  in  $A$ . This follows by an analysis of chords  $W_{q_1} W_{n-1}, W_n W_{q_2-1}, W_{q_1-1} W_n$ , and  $W_{n-1} W_{q_2}$  using the final argument of Case 1. ■

By Lemma 3, it remains for us to count non-degenerate cyclic labelings. We claim that  $C_n(x) = C_n(y)$  iff there is no good fraction between  $x$  and  $y$ . As  $x$  varies, the ordering of points in  $C_n(x)$  changes only when  $C_n(x)$  is degenerate so that two points coincide. It follows that  $C_n(x) = C_n(y)$  when there is no good fraction between  $x$  and  $y$ . If there is a good fraction  $p/q$  with  $x < p/q < y$ , in  $C_n(y)$  there are at least  $p$  integers  $1 \leq i \leq q$  such that  $W_0$  is clockwise of  $W_{i-1}$  and  $W_i$  is clockwise of  $W_0$ , while in  $C_n(x)$  there are fewer than  $p$  such integers. Hence,  $C_n(x)$  and  $C_n(y)$  differ, giving the claim.

We conclude that the number of non-degenerate cyclic labelings is one greater than the number of good fractions in  $(0, 1)$ , hence equal to  $N + 1$ .

This problem was proposed by Alexander Golovanov and Mikhail Ivaniv from Russia.

## Results

The IMO was held in Santa Marta, Colombia, on July 23–24, 2013. There were 527 competitors from 97 countries and regions. On each day contestants were given four and a half hours for three problems.

The top score of 41/42 was shared by Yutao Liu (China) and Eunsoo Jee (South Korea). The USA team won 4 gold and 2 silver medals, placing third behind China and Korea. The students' individual results were as follows.

- Ray Li, who finished 12th grade at Phillips Exeter Academy in Exeter, NH, won a silver medal.
- Mark Sellke, who finished 11th grade at William Henry Harrison High School in West Lafayette, IN, won a gold medal.
- Bobby Shen, who finished 12th grade at Dulles High School in Sugar Land, TX, won a gold medal.
- Thomas Swayze, who finished 12th grade at Canyon Crest Academy in San Diego, CA, won a silver medal.
- James Tao, who finished 11th grade at Illinois Mathematics and Science Academy in Aurora, IL, won a gold medal.
- Victor Wang, who finished 12th grade at Ladue Horton Watkins High School in St. Louis, MO, won a gold medal.